

## Principy anestezie u urologických operací

JONATHAN COUSINS, JEREMY HOWARD a PHILIPPA BORRA

Department of Anaesthesia, Hammersmith Hospital, London, UK

### LITERATURA

- 1 NICE. NICE guideline Pre-op investigations for anaesthetists June 2003, <http://www.nice.org.uk>
- 2 Dodds C, Murray D. Pre-operative assessment of the elderly. *BJA CEPD Reviews* 2001; 1(Issue 6): 181-4
- 3 Nunn JF, Milledge JS, Chen D, Dore C. Respiratory criteria of fitness for surgery and anaesthesia. *Anaesthesia* 1988; 43: 543-51
- 4 Myles PS, Power I, Jamrozik K. Master trial. Epidural block and outcome after major surgery. *Med J Aust* 2002; 177: 536-7
- 5 ACC/AHA. Guidelines for peri-operative cardiovascular evaluation for non-cardiac surgery. Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *JACC* 1996; 27: 910-48
- 6 Mangano DT, Layug EL, Wallace A, Tateo I. Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. *N Engl J Med* 1996; 335: 1713-20
- 7 Poldermans D, Boersma E, Bax JJ *et al.* The effect of bisoprolol on perioperative mortality and myocardial infarction in high-risk patients undergoing vascular surgery. *N Engl J Med* 1999; 341: 1789-94
- 8 Van den Berghe G, Wouters P, Weekers F *et al.* Intensive insulin therapy in critically ill patients. *N Engl J Med* 2001; 345: 1359-67
- 9 Yaster M, Maxwell LG. Paediatric regional anaesthesia. *Anesthesiology* 1989; 70: 324-38
- 10 von Bahr V. Local anaesthesia for inguinal herniorrhaphy. In Eriksson E ed. *Illustrated Handbook in Local Anaesthesia*. Philadelphia: WB Saunders, 1980: 52-4
- 11 Chung FF, Chung A, Meier RH, Lautenschlaeger E, Seyone C. Comparison of perioperative mental function after general anaesthesia and spinal anaesthesia with sedation. *Can J Anesth* 1989; 36: 382
- 12 Sessler DI. Peri-operative heat balance. *Anesthesiology* 2000; 92: 578-96
- 13 Conacher ID, Soormro NA, Rix D. Anaesthesia for laparoscopic urological surgery. *Br J Anaesth* 2004; 93: 859-64
- 14 Goldhill D. Levels of critical care for adult patients. Intensive Care Society UK, 2002; <http://www.ics.ac.uk>
- 15 Tanrikut C, McDougal WS. Acid-base and electrolyte disorders after urinary diversion. *World J Urol* 2004; 22: 168-71
- 16 Jensen V. The TURP syndrome. *Can J Anesth* 1991; 38: 90-6
- 17 Herbert PC, Wells G, Blachman MA *et al.* A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. TRICC Investigators. *N Engl J Med* 1999; 340: 409-17
- 18 NHS. *Better Blood Transfusion*. NHS Executive Clinical Effectiveness Circular, Appendix 1, SC 1998/2, 11 December 1998
- 19 Napier JA, Bruce M, Chapman J *et al.* Guidelines for autologous transfusion II, Peri-operative haemodilution, cell salvage. *Br J Anaesthesia* 1997; 78: 768-71
- 20 Henry DA, Moxey AJ, Carless PA *et al.* Anti-fibrinolytic use for minimising perioperative allogenic blood transfusion. *Cochrane Database Syst Rev* 2001: CD001886
- 21 Chumbley GM, Hall GM, Peter Salmon. Patient controlled analgesia: an assessment by 200 patients. *Anaesthesia* 1998; 53: 216-21
- 22 RCA. *Good practice in the management of continuous epidural analgesia in the hospital setting*. November 2004 Royal College of Anaesthetists UK, <http://www.rcoa.ac.uk/docs/Epid-Analg.pdf>

Korespondence:  
Jonathan Cousins,  
Department of Anaesthesia,  
Hammersmith Hospital,  
London, W12 OHS, UK.  
e-mail: [jc\\_gasman@hotmail.com](mailto:jc_gasman@hotmail.com)